



Phone: 718-443-4000

Fax: 718-443-5000

Sending a Referral to Prime Infusions

Follow the steps below to send a referral to Prime Infusions:

1. Download the desired order form from our website www.primeinfusions.com/referrals
2. Print and fill out all the fields.
3. Fax completed order form with all required documentation listed below to 718-443-5000.

Required Documentation Checklist

If we do not receive all documents below with your referral, the order is subject to delays.

**It may take up to 14 business days for the patient's insurance company to approve or deny our authorization request.*

- ☐ Completed Medication Order Form
- ☐ Patient Demographics
- ☐ Current Medication List and H&P
- ☐ Recent Visit Notes
- ☐ Lab Results
- ☐ Patient's Insurance Card
- ☐ Existing Prior Authorization (if applicable)

How to Use Our Digital Order Forms

Upon downloading the desired form, you will see light blue text box, check box, and circle box fields appear. To fill out the form on your computer, click into these fields to type out the necessary patient, office, clinical history, and therapy administration information. You can copy/paste information from the patient's medical record into this form.

Gather the referring provider's signature to approve the order once you have filled out all fields and send to Prime Infusions via fax at 718-443-5000.

Prescription Requirements

Please be advised that infusion order forms do not serve as valid prescriptions and may not be used as a substitute for one. In accordance with New York State law, a valid prescription must be transmitted electronically or faxed using an official New York State prescription format.

Cerezyme® Referral Form



PATIENT INFORMATION

REFERRAL STATUS: ☐ New Referral ☐ Updated Referral ☐ Referral Renewal

DOB: _____ Patient Name: _____ Patient Phone: _____

Patient Address: _____ Patient Email: _____

☐ NKDA Allergies: _____ Weight (lb/kg): _____ Height: _____

ICD-10 Code (required) _____ ICD-10 Code Description: _____ Last Treatment Date: _____ Last 4 Digits SSN: _____

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City, State, ZIP: _____

Physician Preferred Method of Contact ☐ Fax ☐ Phone ☐ Email: _____

NURSING

☐ Infusion to be administered per Prime Infusions protocols.

LABORATORY ORDERS

☐ CBC ☐ at each dose ☐ every _____

☐ CMP ☐ at each dose ☐ every _____

☐ CRP ☐ at each dose ☐ every _____

☐ Other _____

PREMEDICATIONS

☐ acetaminophen (Tylenol) ☐ 500mg ☐ 650mg ☐ 1000mg ☐ PO ☐ IV

☐ cetirizine (Zyrtec) 10mg ☐ PO ☐ IV

☐ loratadine (Claritin) 10mg ☐ PO ☐ IV

☐ diphenhydramine (Benadryl) ☐ 25mg ☐ 50mg ☐ PO ☐ IV

☐ methylprednisolone (Solu-Medrol) ☐ 40mg ☐ 125mg ☐ PO ☐ IV

☐ hydrocortisone (Solu-Cortef) ☐ 100mg ☐ PO ☐ IV

☐ Other _____

Dose: _____ Route: _____

CEREZYME® THERAPY ADMINISTRATION

☐ 2.5 units/kg three times a week

☐ 60 units/kg once every two weeks

☐ Other: _____

REQUIRED DOCUMENTATION

☐ Patient Demographics

☐ Insurance Card/Information

☐ Progress Notes Supporting DX

☐ Current Medication List and H&P

Consider administering premedication for prophylaxis against infusion reactions and hypersensitivity reactions. **Order is valid for one year unless otherwise noted*

PROVIDER NAME (PRINT)

PROVIDER SIGNATURE

DATE

Cerezyme® Order Form

PHYSICIAN ORDER FORM IS VALID FOR ONE YEAR FROM THE DATE OF SIGNATURE, UNLESS NOTED OTHERWISE.



Phone: 718-443-4000
Fax: 718-443-5000

1. Cerezyme _____ every _____ ☐ days/ ☐ weeks

a) Patient's weight: _____ kg

2. Reconstitution and administration:

- a) Reconstitute each vial of Cerezyme with 10.2 mL of sterile water for injection for a final volume of 10.6 mL. Withdraw only 10 mL from the vial
- b) Dilute reconstituted Cerezyme in a 100 mL bag of Normal Saline for a total volume of 100–200 mL
- c) Run infusion over 2 hours at a rate of 50cc/hr using a sterile, low protein binding filter (size 0.2–1.2 micron), as tolerated

3. Flush IV line with 0.9% Sodium Chloride 10mL prefilled syringe pre/post infusion as needed

4. Premedication only if specified by prescriber:

- a) Diphenhydramine 25mg capsules – take 1–2 capsules 15–30 minutes prior to Cerezyme infusion
- b) Acetaminophen 325mg tablets – take 1–2 tablets by mouth prior to Cerezyme infusion

5. Anaphylaxis kit per protocol:

- a) Epinephrine 0.3mg autoinjector IM every 5 minutes x2 – call EMS
- b) Diphenhydramine 50mg/mL (dispense 1 vial) – administer 50mg IV during anaphylactic reaction
- c) Methylprednisolone 125mg – administer IV push during anaphylactic reaction
- d) 0.9% Sodium Chloride 250mL – administer during anaphylactic reaction

6. Dispense pump and infusion supplies

7. Dispense 8-week supply

- a) Refill: 12-month supply (all above)

8. Nursing orders and plan of treatment:

- a) Nurse to place PIV or use PICC/PORT/CVC to infuse Cerezyme as directed
- b) Assess and monitor vital signs and systems review with each visit
- c) Instruct on the following:
 - i) Disease progress, signs and symptoms, and complications
 - ii) Medication therapy including action, purpose, side effects, and storage of medication and supplies
 - iii) Universal precautions, 911, 24-hour phone number, and when to call RN or physician

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City, State, ZIP: _____

Physician Preferred Method of Contact ☐ Fax ☐ Phone ☐ Email: _____

PROVIDER SIGNATURE

DATE